

A critical look at cosmic ray-cloud relationships



Europska Unija
Ulaganje u budućnost
Projekt je sufinancirala Europska Unija iz
Europskog socijalnog fonda



Jaša Čalogović

Hvar Observatory
Faculty of Geodesy
Zagreb, Croatia




Hvar Observatory

Photosphere and Chromosphere observations



- Group for Solar Physics (6 people):
 - space weather, eruptive processes
 - activity cycle, diff. rotation, convection
 - solar activity & climate
- Group for Stellar Physics (3 people):







10h 20m 53s UT
02.07.2012.
H-alpha
Hvar Solar Telescope

Influence of solar variability on the Earth's climate requires knowledge of

1. Short- and long-term solar variability
2. Solar-terrestrial interactions
3. Mechanisms determining the response of the Earth's climate system to these interactions



Rind, 2002






Solar activity modulates cosmic rays

- Cosmic rays (CR) consist of high-energy particles (mainly protons)
- CR flux of low energy particles is greater than flux of high energy particles ($E^{-\gamma}$)
- Particles with less energy are more influenced by the Sun



“Clear-air” mechanism



Laken & Čalogović, 2015, TOSCA handbook

The hypothesized link between cosmic ray flux and cloud cover


Long-term studies

Svensmark and Friis-Christensen (1997)

- analyzed one solar cycle and reported that global cloud cover changed in phase with the GCR flux by 2-3% → radiative forcing ($0.8 - 1.7 \text{ W/m}^2$) is comparable with greenhouse gases forcing


Marsh and Svensmark, 2000

low clouds (0-3.2km)




Climate sceptics still

use these (incorrect) arguments
Many critics find a forced correlation and heavy debates in the scientific



Long-term cloud data doesn't support GCR-cloud link

Low clouds (<3.2km), global



- Correlation only in low (<3.2km) ISCCP cloud (1983–1995)
- High correlation from 12-month smoothed data (df=4)
- Low (non-significant) correlation from unsmoothed data


Laken, Pallé, Čalogović & Dunne, 2012, SWSC

Artificial correlation b/w low and high cloud

- Measurements are non-cloud penetrating
- Changing number of geostationary → artificial drop in low cloud
- Satellite cloud issues well known (e.g. Hughes, 1984; Minnis, 1989, Tian & Curry, 1989; Rozendall et al. 1995; Loeb & Davies, 1996; Salby & Callaghan, 1997, Campbell, 2004)

Evidence for CR – cloud link is based on low level clouds:


these data are not reliable!



changes in the satellite constellation


Quality of long-term cloud data

Linear ISCCP trend (1983-2008)




Laken, Pallé, Čalogović & Dunne, 2012, SWSC

If linear trends in CR and cloud data are removed correlation becomes weak



Short-term studies - opportunity to test GCR-cloud hypothesis

- Short-term cosmic ray changes comparable to solar cycle amplitude



Čalogović et al., 2010

Advantages:

- Unwanted factors that influence long-term studies removed (ENSO, volcanic eruptions, satellite calibration errors)

Disadvantages:

- Meteorological variability (noise) **increased**
- **Limited** number of high-magnitude Forbush decreases (several per decade)

Short-term studies show conflicting results

- **positive correlations:**

Tinsley & Deen, 1991; Pudovkin & Vertenenko, 1995; Todd & Kniveton, 2001; 2004; Kniveton, 2004; Harrison & Stephenson, 2006; Svensmark *et al.*, 2009; Solovyev & Kozlov, 2009; Harrison & Ambaum, 2010; Harrison *et al.* 2011; Okike & Collier, 2011; Dragić *et al.* 2011; 2013; Svensmark *et al.*, 2012; Zhou *et al.* 2013; Aslam & Badruddin, 2015


- **negative correlations:**

Wang *et al.*, 2006; Troshichev *et al.*, 2008

- **no correlations or inconclusive results:**

Pallé & Butler, 2001; Lam & Rodger, 2002 ; Kristjánsson *et al.*, 2008 ; Sloan & Wolfendale, 2008; Laken *et al.*, 2009; Čalogović *et al.*, 2010; Laken & Kniveton 2011; Laken *et al.*, 2012; Erlykin and Wolfendale, 2013


What is composite?



- Successive averaging of events (in time or space)
- Used to increase signal-to-noise ratio (SNR)
- Enable detection of small amplitude signal against large variability

Event selection affect composites

Example from the literature that used 5 events in its composite:
Individual 5 Fd events plotted against event 2 (19.1.2005)




Laken, Čalogović, Beer and Pallé (2012), *ACPD*

Time period considered matters


MODIS Liquid cloud fraction changes using 5 biggest Fd events

±20 day analysis period



Looking at a short period around the key date can give a false impression of the data.

±100 day analysis period




Examining longer periods shows what 'normal' variations are.

Laken, Čalogović, Beer and Pallé (2012), *ACPD*

Traditional significance tests may not be good enough


Traditional tests (e.g. T/U tests), require minimum sample sizes, specific distributions, and adjustment for autocorrelation



By generating large populations of random events identical in design to a composite with real events, the probability (p) of obtaining a given value by chance in a composite with real events can be accurately known

Laken & Čalogović, SWSC, 2013


Significance testing depends on space and time



Studies using only strong Fd events have usually less than **10** events


Abusing composites: how to make normal changes seem significant

Svensmark et al. 2012, ACPD



Bad approach

Laken & Čalogović, (2013), SWSC



Good approach


Signal detection issues

- **Filtering** - remove irrelevant variations
- **Normalization** - affect magnitude and significance
- **Autocorrelation** - use appropriate statistical tests
- **Signal-to-noise ratios** - affected by area and time period considered

- Weather: **highly variable**, unstable (**non-stationary**), spatio-temporally **autocorrelated**
- No substitute for long datasets: satellite-era data covers three solar cycles.
- ***a posteriori* selection of data**
- Solar—climate links poorly understood
- **Statistical studies**: vulnerable to biased data selection, treatment, assumptions and post-hoc hypotheses.

Conclusions

- **No compelling evidence** to support a global cosmic ray-link using the satellite cloud data (ISCCP, MODIS) with long- or short-term (Fd) studies.
- Satellite cloud data is not suitable for **long-term analysis**
- Co-variance of solar-related parameters (UV, TSI, CR flux, solar wind) make **signal attribution difficult**.
- **Internal variability** at time-scales like the solar cycle complicate signal attribution.
- **Different methodological approaches** produce conflicting results.
- Local effects on cloud can't be dismissed

A sunset over the ocean. The sun is a bright, glowing orb on the left side of the horizon, casting a shimmering reflection on the water. The sky is a deep orange and red, with several wispy, dark clouds scattered across it. The overall scene is serene and dramatic.

Thank you for your attention!

We acknowledge the support of Croatian Science Foundation under the project 6212 „**Solar and Stellar Variability**“ and of European social fund under the project “**PoKRet**”.